

A combination theorem for relatively hyperbolic groups

Emina Alibegović

February 8, 2008

Abstract

In this paper we give new requirements that a tree of δ -hyperbolic spaces has to satisfy in order to be δ -hyperbolic itself. As an application, we give a simple proof that limit groups are relatively hyperbolic.

1 Introduction

In his work on Diophantine equations over free groups Z. Sela introduced limit groups. He showed that this class of groups coincides with the class of ω -residually free groups. Since then few more characterizations of limit groups have been given, as well as structure theorem for limit groups.

This work introduced a lot of interesting questions one might ask about limit groups. We were interested in describing the set of homomorphisms from an arbitrary f.g. group G into a limit group L , $Hom(G, L)$. A key tool in studying $Hom(G, L)$ is a δ -hyperbolic space on which the given limit group L acts freely, by isometries. We construct such a space in Section 3. That the space we constructed is δ -hyperbolic follows from Theorem 2.3 which gives conditions a tree of hyperbolic spaces has to satisfy in order to be a hyperbolic space. The proof of this theorem is given in Section 2 and is an adaptation of the proof of Bestvina-Feighn Combination Theorem to a different setting.

2000 Mathematics Subject Classification. 57M07, 20F65.

Key words and phrases. relatively hyperbolic groups, limit groups.

Existence of such spaces for limit groups gives an answer to the question whether limit groups are hyperbolic relative to their maximal noncyclic abelian subgroups. This question was answered affirmatively by F. Dahmani, [5], who proved a combination theorem for geometrically finite convergence groups using different methods.

Acknowledgements I would like to thank Mladen Bestvina for his support. I will always be grateful for the time and knowledge he generously shared with me.

2 A combination theorem

Let X be a connected finite cell complex which is a graph of spaces. There is a map $p : X \rightarrow \Gamma$ onto a finite graph Γ . Let X_e denote the preimage of a midpoint of an edge e in Γ , and let X_v denote the preimage of a component of $\Gamma \setminus \{\text{midpoints of all edges}\}$ that contains the vertex v . We require that X_e and X_v are connected and that their inclusions into X induce inclusions on fundamental groups. There is an induced map $\tilde{p} : \tilde{X} \rightarrow T$ from the universal cover of X onto a $\pi_1(X)$ -tree T so that $T/\pi_1(X)$ is isomorphic to Γ . We say that X is a *graph of negatively curved spaces* if every vertex space X_v is negatively curved. As a reminder a cell complex X is said to be negatively curved if there exists a constant $A = A(X)$ so that each inessential circuit bounds a disk of combinatorial area which is bounded above by A times the combinatorial length of the circuit (we assign length 1 to each edge).

We know that in δ -hyperbolic spaces geodesic triangles are δ -thin. We have a similar fact for polygons, in fact for quasigeodesic polygons. The following proposition can be found in [6] and [1].

Proposition 2.1. *Let Z be a δ -hyperbolic space and let $\tau \geq 1$ be a constant. There is a function $B(x) = O(\log x)$ and a linear function $C(x)$ each depending only on Z and τ with the following property. If $\Delta : D^2 \rightarrow Z$ is a disk with boundary a k -sided τ -quasigeodesic polygon, then there is a finite tree S and a map $r : D^2 \rightarrow S$ such that:*

1. *the number of valence one vertices of S is k ,*
2. *for a and b in S^1 , $d_Z(\Delta(a), \Delta(b)) \leq d_S(r(a), r(b)) + B(k)$,*
3. *$r^{-1}(s)$ is a properly embedded finite tree in D^2 for $s \in S$,*

4. if E is an edge of S , then r restricted to $r^{-1}(\text{Interior}(E))$ is an I -bundle,
5. for a_1, b_1 (respectively a_2, b_2) in the same side of the polygon and satisfying $r(a_1) = r(a_2) \in E$ and $r(b_1) = r(b_2) \in E$, we have

$$\ell(\Delta(\text{the circular arc } a_1b_1 \text{ in the edge of the polygon})) \leq$$

$$C(\ell(\Delta(\text{the circular arc } a_2b_2 \text{ in the edge of the polygon}))).$$

Such a map r is called a resolution of the quasigeodesic polygon. A singular fiber of the resolution is a fiber which is not isomorphic to I .

Partial qi-embedded condition We will say that a graph of spaces X satisfies the partial quasiisometrically embedded condition if every edge space \widetilde{X}_e is quasiisometrically embedded in at least one of the vertex spaces \widetilde{X}_v and \widetilde{X}_w , where v and w are endpoints of the edge e in Γ . We further ask that all qi-constants be equal.

Qi-consistency condition A graph of groups X satisfies the qi-consistency condition if the following holds: If one of the edge spaces adjacent to a vertex space \widetilde{X}_v qi-embeds into it, then the same is true for all adjacent edge spaces. We will call such vertex spaces *good*.

Compact intersection condition A graph of spaces X satisfies the compact intersection condition if whenever the edge spaces \widetilde{X}_e and \widetilde{X}_f qi-embed into the same vertex space \widetilde{X}_v then the intersection of any of their Hausdorff neighborhoods in \widetilde{X}_v is a compact set.

Remark 2.2. The compact intersection condition gives us the following. Suppose that \widetilde{X}_e and \widetilde{X}_f qi-embed into the same vertex space \widetilde{X}_v . If we fix a constant $k < \infty$, then

$$L = \max\{\ell(S_1) : \exists S_2 \subset \widetilde{X}_f \text{ so that } d_H(S_1, S_2) \leq k\}$$

is finite, where S_1, S_2 are quasigeodesics in \widetilde{X}_e and \widetilde{X}_f , respectively. Note that L depends on the qi-constants for S_1 and S_2 .

Theorem 2.3. *If a tree of negatively curved spaces X satisfies the partial qi-embedded, the qi-consistency and the compact intersection conditions, then X is negatively curved.*

We would like to show that X satisfies subquadratic isoperimetric inequality, which will then imply that X is a hyperbolic space ([2], [6]). We

will use the techniques employed by Bestvina and Feighn in the proof of their Combination theorem ([1]).

Let $\gamma : S^1 \rightarrow X$ be a circuit that is transverse to and has nonempty intersection with $\cup\{X_e : e \text{ edge of } T\}$. We may also assume that γ is contained in the 1-skeleton of X ([4]). Following [1] we talk about *good disks*. There is a disk $\Delta : D^2 \rightarrow X$ with boundary γ . The set $\mathcal{W} = \Delta^{-1}(\cup\{X_e : e \text{ edge of } T\})$ divides D^2 into regions that are mapped into negatively curved vertex spaces, see Figure 1. Elements of \mathcal{W} are called *walls*. We may assume that Δ has the following properties:

- (1) The set \mathcal{W} consists of properly embedded arcs in D^2 .
- (2) The length of $\Delta(\cup\mathcal{W})$ in X is minimal over all disks satisfying (1).
- (3) The closures of the components of $\Delta(D^2 \setminus (\cup\mathcal{W}))$ have areas bounded by A times the length of their boundaries, where A is a constant.
- (4) Define \mathcal{L} to be the set of closures of the components of $S^1 \setminus (S^1 \cap \cup\mathcal{W})$. We may assume that γ restricted to each element of \mathcal{L} is a geodesic in the appropriate X_v . We view γ as a polygon whose sides are elements of \mathcal{L} . Hence the number of sides of γ can be no more than $\ell(\gamma)$ (the length of each side is at least 1).

A disk is good if it satisfies (1)-(4).




Figure 1: Decomposition by walls of a disk Δ into polygons

Our goal is to bound the area of Δ by a subquadratic function of the length of its boundary. For a good disk Δ , by property (3), we have

$$\text{Area}(\Delta) \leq A(2\ell(\Delta(\cup \mathcal{W})) + \ell(\Delta(\cup \mathcal{L}))) \leq A(2\ell(\Delta(\cup \mathcal{W})) + \ell(\gamma)).$$

Therefore we need to bound $\ell(\Delta(\cup \mathcal{W}))$ in terms of $\ell(\gamma)$.

Let us denote by \mathcal{P} the set of closures of the components of $D^2 \setminus \cup \mathcal{W}$, and let P be an element of \mathcal{P} . If $\Delta(P)$ is contained in a good vertex space X_v , then the map Δ restricted to ∂P is a τ -quasigeodesic polygon in X_v . Note that each wall $W \in \mathcal{W}$ is a side of at least one polygon $P \in \mathcal{P}$ for which $\Delta(P)$ is contained in a good vertex space, and so we only need to consider such polygons. In what follows in order to avoid a cumbersome notation we will write $\ell(W)$ and P when we really mean $\ell(\Delta(W))$ and $\Delta(P)$.

Claim 1:

$$\sum \{\ell(W) : W \subset \text{a bigon } P \subset X_v\} \leq \tau \ell(\gamma).$$

Proof of Claim If P is a bigon whose one side is a wall W , then the other side s of that bigon is an element of \mathcal{L} . According to (4) in the definition of good disks the images of the elements of \mathcal{L} under Δ are geodesics in appropriate vertex spaces and hence $\ell(W) \leq \tau \ell(s)$. The claim follows.

Claim 2:

$$\sum \{\ell(W) : W \subset \text{an } m\text{-gon } P, m \geq 4\} = O(\ell(\gamma) \log(\ell(\gamma))).$$

Proof of Claim A polygon P is a τ -quasigeodesic polygon and can be resolved using Lemma 2.1. We call a point $w \in W$ a singular point if it lies in a singular fiber of the resolution of the polygon P . These points will decompose W into the union of closed segments V . Let $\mathcal{V}(W)$ denote the set of all such V 's, and let $\mathcal{V} = \cup \{\mathcal{V}(W) : W \in \mathcal{W}\}$. Since

$$\sum \{\ell(W) : W \subset \text{an } m\text{-gon } P, m \geq 4\} = \sum \{\ell(V) : V \in \mathcal{V}\}$$

we need to bound $\ell(V)$, for all $V \in \mathcal{V}$.

Note that singular fibers decompose the polygon P into the union of quadrilaterals and triangles. The case of triangles is relatively easy to handle. One of the sides of this triangle is some $V \in \mathcal{V}$ and another, call it S , is contained in S^1 . Considering how the resolution was formed, we have that

$\ell(V) \leq C(\ell(S))$, where the function C is a linear function from Proposition 2.1 (5). Hence,

$$\sum \{\ell(V) : V \in \mathcal{V}\} \leq D(\ell(\gamma)) + \sum \{\ell(V) : V \text{ side of quadrilateral in } P\}$$

where D is a linear function. Let us consider the case of a quasigeodesic quadrilateral Q with sides V_1 and V_2 (contained in X_e and X_f , respectively) that are joined by singular fibers of the resolution r of the polygon P . We may assume that V_1 is shorter of the two. According to Proposition 2.1 the distance between the images under Δ of the two endpoints of a fiber is at most B (the number of sides of P) $\leq B(\ell(\gamma))$. Since Q is contained in δ -hyperbolic space, it is not hard to show that as long as $\ell(V_1) > 2\tau B(\ell(\gamma)) + d$, where $d = d(\tau, \epsilon, R)$, we can find subsegments $S_1 \subset V_1$ and $S_2 \subset V_2$ such that $d_H(S_1, S_2) \leq 12\delta + 4R(\tau + 1) + 2\epsilon$. If $\ell(S_1) > L$, (L from Remark 2.2, where the constant k is $12\delta + 4R(\tau + 1) + 2\epsilon$), then both S_1 and S_2 belong to X_e . By doing a surgery, see Figure 2, we can shorten the length of the walls in our disk, which contradicts the assumption that Δ was a good disk.

Thus, if

$$\ell(V) > 2\tau B(\ell(\gamma)) + d + L, \text{ for some } V \in \mathcal{V},$$

Δ is not a good disk. To finish the proof we need to know what the cardinality of \mathcal{V} is. The number of $V \in \mathcal{V}$ is proportional to the number of singular fibers inside Δ . On the other hand, there can be no more singular fibers than there are triangles in the triangulation of the polygon γ . Hence, $\text{cardinality}(\mathcal{V}) = O(\ell(\gamma))$, and our claim follows.

Proof of Theorem 2.3. If Δ is a good disk we already noticed that

$$\begin{aligned} \text{Area}(\Delta) &\leq A(2\ell(\Delta(\cup \mathcal{W})) + \ell(\Delta(\cup \mathcal{L}))) \\ &\leq A(2\ell(\Delta(\cup \mathcal{W})) + \ell(\gamma)) \\ &\leq 2A(\tau\ell(\gamma) + O(\ell(\gamma)\log(\ell(\gamma)))) + A\ell(\gamma) \\ &= O(\ell(\gamma)\log(\ell(\gamma))) \end{aligned}$$

The last inequality follows from Claims 1 and 2. Therefore, X satisfies sub-quadratic isoperimetric inequality, and our proof is finished. \square

3 Application to limit groups

The goal of this section is to produce a δ -hyperbolic space on which a given limit group acts freely, by isometries. We will in fact consider a slightly




Figure 2: Surgery: the lengths of the central 'parallel' curves are much shorter than the lengths of walls connecting them. We change a disk by pushing this tunnel down into X_c and get a new disk: a good disk. For the better visibility we drew X_e as two dimensional.

larger class of groups, \mathcal{C} . We describe elements of \mathcal{C} (\mathcal{C} -groups, for short) inductively.

Definition 3.1. A torsion-free, f.g. group G is a depth 0 \mathcal{C} -group if it is either an f.g. free group, or an f.g. free abelian group or the fundamental group of a closed hyperbolic surface. A torsion-free f.g. group G is a \mathcal{C} -group of depth $\leq n$ if it has a graph of groups decomposition with three types of vertices: abelian, surface or depth $\leq (n - 1)$, cyclic edge stabilizers and the following holds:

- Every edge is adjacent to at most one abelian vertex v . Further, G_v , the stabilizer of v , is a maximal abelian subgroup of G .
- Each surface vertex group is the fundamental group of a surface with

boundary, and to each boundary component corresponds an edge of this decomposition. Each edge group is conjugate to a boundary component.

- The stabilizer of a depth $\leq (n - 1)$ vertex v , G_v , is \mathcal{C} -group of depth $\leq (n - 1)$. The images in G_v of incident edge groups are distinct maximal abelian subgroups of G_v (i.e., cyclic subgroups generated by distinct, primitive, hyperbolic elements of G_v).

We say that the depth of a \mathcal{C} -group G is the smallest n for which G is of depth $\leq n$.

We will get a hyperbolic space on which a \mathcal{C} -group G acts freely by induction on its depth. If G is a depth 0 \mathcal{C} -group, we take a tree, a horoball or \mathbb{H}^2 .

Let G be a depth n \mathcal{C} -group. For the vertex groups in the decomposition of G as above we have the desired spaces by induction. Let X/G be a graph of spaces corresponding to this splitting, and X its universal cover. Let T_G be the corresponding graph of groups, and let T be a tree so that $T/G = T_G$. Our goal is to show that X satisfies the hypotheses of Theorem 2.3 and consequently that X is a hyperbolic space.

The requirement we imposed on the splitting of G guarantees that X/G satisfies the partial qi-embedded condition and the qi-consistency condition. Namely, the generators of all edge groups adjacent to a nonabelian vertex group are identified with hyperbolic elements of that vertex group, hence the corresponding edge spaces in X are glued along quasigeodesics in the relevant vertex space. Just as a note, no edge space qi-embeds into a vertex space which is a horoball.

Claim: X/G satisfies the compact intersection condition.

Proof of Claim: Suppose edge spaces \widetilde{X}_e and \widetilde{X}_f qi-embed into the vertex space \widetilde{X}_v . We noticed that they are glued along quasigeodesics, say c_1 and c_2 respectively. There are elements g_1 and g_2 of $\pi_1 X_v$ that act as translations along c_1 and c_2 , respectively. If the Hausdorff distance between quasigeodesics c_1 and c_2 is bounded, we conclude that g_1 and g_2 fix the same two points in $\partial \widetilde{X}_v$. Hence, they are both contained in a unique elementary group that is virtually cyclic. Since we have no torsion elements, this elementary group is cyclic, contradicting the choice of splitting for G . \square

The definition of relatively hyperbolic groups appears in many forms. We use the one given by M. Gromov in [6], 8.6.

Let X be a complete hyperbolic locally compact geodesic space with a discrete free isometric action of a group Γ such that the quotient $V = X/\Gamma$ is quasiisometric to the union of k copies of $[0, \infty)$ joined at 0. Lift k rays that correspond to ∂V to rays $r_i : [0, \infty) \rightarrow X$, $i = 1, \dots, k$. Let h_i be the horofunction corresponding to r_i and let $r_i(\infty)$ be the limit point of r_i . Denote by $\Gamma_i < \Gamma$ the stabilizer of $r_i(\infty)$ and assume that it preserves h_i . Denote by $B_i(\rho)$ the horoballs $h_i^{-1}(-\infty, \rho) \subset X$ and assume that for sufficiently small ρ the intersections $\gamma B_i(\rho) \cap B_j(\rho)$ is empty unless $i = j$ and $\gamma \in \Gamma_i$. Let

$$\Gamma B(\rho) = \bigcup_{i, \gamma} \gamma B_i(\rho),$$

$i = 1, \dots, k$, $\gamma \in \Gamma$. Let $X(\rho) = X \setminus \Gamma B(\rho)$, and assume that $X(\rho)/\Gamma$ is compact for all $\rho \in (-\infty, \infty)$.

Definition 3.2. We say that a group Γ is hyperbolic relative to the subgroups $\Gamma_1, \dots, \Gamma_k$ if Γ admits an action on some X as above, and where Γ_i are the stabilizers of h_i .

After inspection of the action of $G \in \mathcal{C}$ on the δ -hyperbolic space X that we constructed above, we see that all the requirements of Definition 3.2 are satisfied. Hence, we have proved:

Theorem 3.3. *Groups in \mathcal{C} are hyperbolic relative to the collection of the conjugacy classes of their maximal noncyclic abelian subgroups.*

Since limit groups belong to the class \mathcal{C} , see [8] Theorem 3.2. and Theorem 4.1., the consequence of this theorem is the following corollary:

Corollary 3.4. *Limit group L is hyperbolic relative to the collection of representatives of conjugacy classes of its maximal noncyclic abelian subgroups.*

This fact was also noted in the work of F. Dahmani, [5].

Several nice properties follow from the relative hyperbolicity. I. Bumagin showed that the conjugacy problem is solvable for a group G which is hyperbolic relative to a subgroup H with solvable conjugacy problem; hence limit groups have solvable conjugacy problem ([3]). D. Y. Rebecchi showed that a group G hyperbolic relative to a biautomatic subgroup H is itself biautomatic, [7]. We therefore conclude that limit groups are biautomatic.

References

- [1] M. BESTVINA AND M. FEIGN, *A combination theorem for negatively curved groups*, J. Differential Geom., 35 (1992), pp. 85–101.
- [2] B. BOWDITCH, *A short proof that a subquadratic linear isoperimetric inequality implies a linear one*, Michigan Math. J., 42 (1985), pp. 103–107.
- [3] I. BUMAGIN, *The conjugacy problem for relatively hyperbolic groups*. preprint, <http://front.math.ucdavis.edu/math.GR/0308171>, 2003.
- [4] J. BURILLO AND J. TABACK, *Equivalence of geometric and combinatorial Dehn functions*. to appear, 2002.
- [5] F. DAHMANI, *Combination of convergence groups*. preprint, <http://www-irma.u-strasbg.fr/irma/publications/2002/02018.shtml>, 2002.
- [6] M. GROMOV, *Hyperbolic groups*, in Essays in group theory, S. Gersten, ed., Springer-Verlag, 1987, pp. 75–264.
- [7] D. Y. REBBECHI, *Algorithmic properties of relatively hyperbolic groups*. thesis, <http://front.math.ucdavis.edu/math.GR/0302245>, 2001.
- [8] Z. SELA, *Diophantine geometry over groups I: Makanin-Razborov diagrams*, IHES Publ. Math., 93 (2001), pp. 31–105.